

B
IV
HAD

Science Reports of the Tōhoku Imperial University
Biology, Vol. 7, No. 4, 1932.

REPORT OF THE BIOLOGICAL SURVEY OF MUTSU BAY.

24. THE PELAGIC CILIATA, SUBORDER TINTINNOINEA¹.

226 :

By

YOSHINE HADA.

The Akkeshi Marine Biological Station of the Hokkaido University, Akkeshi, Hokkaido, Japan.

(With 26 text-figures)

(Received October 3, 1932)

INTRODUCTION

The present work is based upon the materials obtained by myself in surface collections and vertical hauls during August, 1929-September, 1930 at numerous stations scattered throughout Mutsu Bay, and upon the plankton collected with a surface tow net twice a month during 1927-1931 and preserved in the Asamushi Marine Biological Station of the Tōhoku Imperial University.

In this paper I have recorded 34 species belonging to 12 genera and 8 families according to KOFORD and CAMPBELL's systematic arrangement of the Tintinnoinea (1929). Of these, 7 seem to be new to science.

It is my pleasure and duty to offer my sincere acknowledgment to Professor Dr. C. A. KOFORD and Dr. A. S. CAMPBELL, of the Zoological Department of the University of California, for their kind advice and help in identification of the species and in the accomplishment of this investigation. In collection and examination of the materials I am indebted to Dr. S. KOKUBO and Mr. T. TAMURA, of the Asamushi Marine Biological Station.

SYSTEMATIC PART

Class CILIATA.


Order HETEROTRICHIDA.

Suborder TINTINNOINEA.

Key to genera.

1. Lorica consisting of a bowl with or without an aboral horn; wall

¹) Contributions from the Marine Biological Station, Asamushi, Aomori-Ken. No. 93.



composed of a fine primary structure and an agglomerated material.

Genus *Tintinnopsis*.

2. Lorica with a very low, hyaline collar and a coarsely constituted bowl.

Genus *Stenosemella*.

3. Lorica consisting of a higher, hyaline, subcylindrical collar with spiral turns and a coarser bowl with an alveolar pattern or agglomerated particles.

Genus *Codonellopsis*.

4. Lorica usually bell-shaped, with a flaring collar divided by a nuchal constriction from a bowl; wall bilamellate, with an irregularly polygonal reticulation.

Genus *Cyttaroclysis*.

5. Lorica chalice-shaped; wall composed of separated lamellae with fine, subuniform, hexagons.

Genus *Parafavella*.

6. Lorica bell-shaped, having two elevated bands; wall weakly bilaminated; surface with minute irregular polygons or fine plications.

Genus *Psychoclysis*.

7. Lorica with double collars, inner one a little higher than the outer denticulated with short triangular teeth; wall bilamellate, usually hyaline, but not structureless.

Genus *Acanthostomella*.

8. Lorica elongated chalice-shaped; aboral end drawn out into a simple caudal lance; wall generally hyaline, composed of almost separated laminae fused partially in the aboral end.

Genus *Parundella*.

9. Lorica vase- or cup-shaped, with a fairly developed inner collar by the reason of the thickened wall in the suboral region; wall trilamellate, translucent.

Genus *Proplectella*.

10. Lorica consisting of a low funnel-shaped collar and an elongated bowl with a few ridges or lines; oral rim entire and circular; wall translucent.

Genus *Amphorella*.

11. Lorica subcylindrical, open at the both ends; wall typically hyaline.

Genus *Tintinnus*.

12. Lorica elongate, tubular; collar an inverted, truncate, conical cone; shaft subcylindrical, with some fins at the aboral region; aboral end usually open; wall almost hyaline.

Genus *Salpingella*.

Family *Codonellidae*.

Genus *TINTINNOPTYSIS* STEIN, 1867.

1. *Tintinnopsis beroidea* STEIN.

Tintinnopsis beroidea STEIN, * 1867; Koroid and CAMPBELL, 1929, p. 28, fig. 26; HADA,

* Indicates literature which I have not examined.

1932 b, p. 41, fig. 2.

Length, 61(50-74) μ ; oral diameter, 35(31-40) μ .
Occurs in February-April and September; common.

2. *Tintinnopsis urnula* MEUNIER.
Tintinnopsis tubulosoides MEUNIER, 1910, p. 145, pl. 13, figs. 21-25; KOROID and CAMPBELL, 1929, p. 50, fig. 20; HADA, 1932 b, p. 42, fig. 3.

Length, 60 μ ; oral diameter, 40 μ .
Occurs in March and April; very rare.

3. *Tintinnopsis tubulosoides* MEUNIER.
Tintinnopsis tubulosoides MEUNIER, 1910, p. 139, pl. 12, figs. 10, 11; KOROID and CAMPBELL, 1929, p. 49, fig. 74; HADA, 1932 b, p. 43, fig. 5.

Length, 94(83-104) μ ; oral diameter, 36(34-40) μ .
Occurs in April; rare.

4. *Tintinnopsis tenuis*, n. sp.
Text-figure 1.
Differs from *Tintinnopsis acuminata* DADAY and *Tps. beroidea* STEIN in the rounded aboral end and in the presence of the faint spiral structure, from *Tps. kargiacensis* BRANDT and *Tps. rotundata* JØRGENSEN in dimensions and in a sparse agglomerated material, and from *Tps. minuta* WAMES in being larger and in more slender proportions.

Length, 60(54-64) μ ; oral diameter, 27(25-29) μ .
Occurs in May-October; rare.


Lorica elongated capsular, 2.0-2.5 oral diameters in length; oral rim usually entire; bowl cylindrical; aboral end generally hemispherical; wall thin, subuniform, 0.03 of the oral diameter in thickness, showing a slight spiral structure in the suboral part, with fine and sparse agglomerations. Length, 60(54-64) μ ; oral diameter, 27(25-29) μ .
Occurs in May-October; rare.

Differs from *Tintinnopsis acuminata* DADAY and *Tps. beroidea* STEIN in the rounded aboral end and in the presence of the faint spiral structure, from *Tps. kargiacensis* BRANDT and *Tps. rotundata* JØRGENSEN in dimensions and in a sparse agglomerated material, and from *Tps. minuta* WAMES in being larger and in more slender proportions.

5. *Tintinnopsis congregata*, n. sp.
Text-figure 2.

Lorica ovoidal, 2.3 oral diameters in length, consisting of a short subcylindrical collar and an ovate bowl, widest near the middle, its greatest transdiameter 1.8 of the oral diameter; oral rim ragged: shoulder gradually sloping; aboral end rounded; wall rather coarse and thick, without a spiral structure.


TINTINNOINEA OF MUTSU BAY



Text-fig. 1. *Tintinnopsis tenuis*, n. sp. $\times 750$.

Text-fig. 2. *Tintinnopsis congregata*, n. sp. $\times 750$.

Text-fig. 3. *Tintinnopsis lohmanni* LAACKMANN. $\times 750$.



7. *Tintinnopsis directa*, n. sp.

Text-figure 4.

Tintinnopsis sp. (*T. campanula* var.?) OKAMURA, 1907, p. 139, pl. 6, fig. 6¹.
Tintinnopsis patula (part), KOFOID and CAMPBELL, 1929, p. 43.

Lorica tall campanulate, 1.6–2.2 oral diameters in length; oral rim irregular, flaring (60°–92°); suboral region somewhat tapering, conical (5°–10°), laid up with about 6 spiral turns, narrowest at the basal portion of the subcylindrical part, its smallest transdiameter 0.68–0.82 of the oral diameter; posterior region subspherical, with a rounded aboral end, 0.80–0.95 oral diameters in transdiameter; wall rather coarse in the posterior part, about 0.035 diameters in thickness at the thickest portion of the aboral region.

Length, 46(44–48) μ ; oral diameter, 20 μ .

Occurs in February–April; rare.

Differs from *Tintinnopsis compressa* DADAY in the presence of the marked subcylindrical nuchal part and from *Tps. nucula* (FOL) in the widened bowl, in the hemispherical aboral end and in the less shouldered collar.

6. *Tintinnopsis lohmanni* LAACKMANN.

Text-figure 3.

Tintinnopsis lohmanni LAACKMANN, 1906, p. 20, pl. 1, figs. 10, 11, pl. 2, fig. 23.

Tintinnopsis sp. BRANDT, 1906, pl. 17, figs. 1, 3; 1907, p. 180.

Tintinnopsis tubulosa, var. *lohmanni*, JÖRGENSEN, 1927, p. 7.

Tintinnopsis subacuta (part), KOFOID and CAMPBELL, 1929, p. 47.

Lorica stout flask-shaped, 2 oral diameters in length, cylindrical anteriorly, enlarged aborally; oral rim more or less irregular, no oral flare; tubular part 0.40–0.43 of the total length in length, provided with 2 oral turns; aboral region subspherical or broadly conical, 1.14–1.23 oral diameters in transdiameter at the posterior 0.3 of the total length; wall agglomerated rather coarsely, 0.08 oral diameters in thickness.

Length, 60 μ ; oral diameter, 30 μ .

Occurs in August; rare.

Differs from *Tintinnopsis compressa* DADAY in contour being composed of the oral cylindrical part and the expanding aboral region, from *Tps. subacuta* JÖRGENSEN in the shortened anterior tubular part and in the shape of the aboral end, and from *Tps. turgida* KOFOID and CAMPBELL in the short suboral region and in the presence of the spiral structure in the same portion.

8. *Tintinnopsis bütschlii* DADAY.

Text-figure 5.

Tintinnopsis Bütschlii DADAY, 1887, p. 556, pl. 20, figs. 4, 5; KOFOID and CAMPBELL, 1929, p. 29, fig. 85.

Tintinnopsis campanula var. *bütschlii* (part), BRANDT, 1907, p. 151.

Lorica campanulate, consisting of a broadly expanding and evinated oral region and a convex conical bowl, its length 0.94–1.05 oral diameter; oral rim roughened, conical (about 130°); bowl narrowest at the upper third of the lorica, its least transdiameter 0.41–0.45 of the oral diameter, dilated posteriorly a little (7°–9°), 0.43–0.48 of the oral diameter in greatest transdiameter at the posterior 0.25 of the total length; aboral end hemispherical; wall 0.023–0.027 oral diameters in thickness, with a trace of a spiral structure in the suboral nuchal region.


Length, 88(84–92) μ ; oral diameter, 88(80–92) μ ; greatest transdiameter of the bowl, 40(38–43) μ .

Occurs in August–October; common.

Differs from *Tintinnopsis cyathus* DADAY and *Tps. everta* KOFOID and CAMPBELL in the more spreading oral flare and from *Tps. mortensenii* SCHMIDT in the shape of the oral flare which is not so much everted in this species as in the last.

9. *Tintinnopsis karajacensis* BRANDT.

Text-figure 6.



Tintinnopsis karajacensis BRANDT, 1896, p. 57, pl. 3, fig. 5; 1906 (part), 19, figs. 5, 10, 12, pl. 26, fig. 3; 1907, p. 162; LAACKMANN, 1906, p. 21, pl. 1, figs. 12-14; JÖRGENSEN, 1927, pp. 5, 7; KOROID and CAMPBELL, 1929, p. 37, fig. 38.

Lorica cylindrical, 2.0-2.7 oral diameters in length, oral rim ragged; aboral end rounded or disfigured as the result of irregularly agglomerated particles; wall coarse, having several slight spiral turns in the anterior half.

fig. 4; 1900b, p. 18; BRANDT, 1906, pl. 19, fig. 3, pl. 24, fig. 16, pl. 26, figs. 7, 8; 1907, p. 160; OKAMURA, 1907, p. 137, pl. 6, fig. 56; ENTZ, JR. (part), 1909, pl. 9, fig. 2, pl. 12, fig. 4, pl. 21, fig. 6; MERKLE, 1909, p. 153, pl. 2, figs. 13, 24; MEUNIER, 1910, p. 138, pl. 12, figs. 5-9; JÖRGENSEN, 1927, pp. 5, 7; KOFOID and CAMPBELL, 1929, p. 38, fig. 95.

Tintinnopsis radix forma *subrotundata* LAACKMANN, 1913, p. 23, pl. 2, fig. 32.

Tintinnopsis radix forma *carta-subrotundata* LAACKMANN, 1913, p. 23, pl. 2, fig. 34.

Lorica elongate, tubular, usually straight, 4.5 oral diameters in length; oral rim ragged; aboral end rounded or shaped somewhat irregularly; wall agglomerated roughly, but comparatively thin, 0.04 of the oral diameter in thickness, without a spiral structure.

Length, 151 μ ; oral diameter, 34 μ .

Occurs in September; very rare.

Differs from *Tintinnopsis karajacensis* BRANDT in the longer lorica and in slender proportions and from *Tps. cochleata* (BRANDT) in lack of the spiral structure.


11. *Tintinnopsis tocantinensis* KOFOID and CAMPBELL.

Text-figure 8.

Tintinnopsis aperta var. a BRANDT, 1906, pl. 25, figs. 2, 7; 1907, pp. 129, 177.
Tintinnopsis tocantinensis KOFOID and CAMPBELL, 1929, p. 48, fig. 46.



- Text-fig. 4. *Tintinnopsis directa*, n. sp. $\times 650$.
Length, 111-172 μ ; oral diameter, 55-64 μ .
Occurs in June and July; rare.
- Differs from *Tintinnopsis cochleata* (BRANDT) in less extensive spiral organization and in roughened agglomeration, from *Tps. lobiancoi* DADAY in the shorter lorica, and from *Tps. rotundata* JÖRGENSEN in more slender proportions and in the shape of the aboral end.
10. *Tintinnopsis lobiancoi* DADAY.
Text-figure 7.
- Tintinnopsis lobiancoi* DADAY, 1887, pp. 545, 553, pl. 19, fig. 27; CLEVE, 1900a, p. 17,
Tintinnopsis lobiancoi DADAY $\times 400$. CAMPBELL $\times 600$.



- Text-fig. 8. *Tintinnopsis tocantinensis* KOFOID and CAMPBELL $\times 400$.
Text-fig. 9. *Tintinnopsis kofoidi* HADA $\times 350$.

Lorica elongated, 4.7 oral diameters in length, cylindrical anteriorly, expanding posteriorly, tapering distally into a stout aboral horn; dilated part not spiraled, 1.2 of the oral diameter in transdiameter; aboral horn conical (35°), obliquely or irregularly open at the tip; wall thick and coarse.

Length, $103\ \mu$; oral diameter, $22\ \mu$.

Occurs in September; very rare.

Differs from *Tintinnopsis aperta* BRANDT in the absence of the spiral structure at the enlarged region and in having the stout aboral horn.

12. *Tintinnopsis kofoidi* HADA.

Text-figure 9.

Tintinnopsis kofoidi HADA, 1932 a, p. 210, figs. 2, 3; 1932 b, p. 44, fig. 6.

Occurs in July; rare.

13. *Tintinnopsis radix* (IMHOFF) BRANDT.

Text-figure 10.

Codonella radix IMHOFF, 1886, p. 103.
Tintinnopsis Davidoffi DADAY, 1887, p. 552, pl. 19,

fig. 23.
Tintinnopsis Davidoffi var. *cylindrica* (part), DADAY, 1887, p. 553, pl. 19, fig. 25.

Tintinnopsis Davidoffi var. *longicauda*, DADAY, 1887, pp. 545, 553, pl. 19, fig. 26.
Tintinnopsis curvicauda DADAY, 1887, p. 554, pl. 19, fig. 33.

Tintinnopsis fracta BRANDT, 1906 pl. 23, figs. 1, 3-5, 9-13, pl. 31, fig. 8; 1907, p. 174; OKAMURA, 1907, p. 137, pl. 6, fig. 57.


Tintinnopsis radix BRANDT, 1907, p. 20; LAACKMANN, 1913, p. 17, pl. 2, figs. 17-20, 27-28; KOFOD and CAMPBELL, 1929, p. 45, fig. 93.

Tintinnopsis radix forma *typica* LAACKMANN, 1913, p. 22.

Tintinnopsis radix forma *curta* LAACKMANN, 1913, p. 23, pl. 2, figs. 21-24, 26.

Tintinnopsis radix forma *cylindrica*, LAACKMANN, 1913, p. 23, pl. 2, figs. 25, 29-31.

Lorica elongate, slender, tubular, 6.0-9.5 oral diameters in length; oral rim generally



Family Codonellopsidae.

Genus STENOSEMELLA JÖRGENSEN, 1924.

14. *Stenosemella nivalis* (MEUNIER) KOFOID and CAMPBELL.

Text-figure 11.

Codonella ventricosa, ENTZ, SR., 1884, p. 413, pl. 24, fig. 24.

Tintinnopsis ventricosa, DADAY, 1887, pp. 546, 559, pl. 20, figs. 19, 20.

Tintinnopsis nivalis (part), LAACKMANN, 1906, p. 19, pl. 1, fig. 4, pl. 3, figs. 48-50; CAMP-

BELL, 1926, pp. 173-236, pl. 12-15, text-figs. A-C.

Tintinnopsis nivalis MEUNIER, 1910, p. 143, pl. 13, figs. 26, 27.

Stenosemella nucula, JÖRGENSEN, 1927, p. 8, fig. 7.

Stenosemella nivalis, KOFOID and CAMPBELL, 1929, p. 69, fig. 136.

Lorica ovoidal, 1.8-2.2 oral diameters in length; collar somewhat concave conical, 0.08-0.10 of the total length in height; bowl widest a little upper the middle of the lorica, 1.9 oral diameters in transdiameter; aboral region subacute; wall of the collar thin and hyaline, about 0.05 of the oral diameter in thickness, sometimes with a few foreign particles, wall of the bowl with a rather coarse agglomerated material.

Length, $43(40-44)\ \mu$; oral diameter, $21(20-22)\ \mu$; transdiameter of the bowl, $39(36-41)\ \mu$.

Occurs throughout the year; rare.

Differs from *Stenosemella pacifica* KOFOID and CAMPBELL in lack of fenestrae at the base of the collar and from *S. ventricosa* (CLAPARÈDE LACHMANN) in dimensions.

Genus CODONELLOPSIS JÖRGENSEN, 1924.

15. *Codonellopsis pusilla* (CLEVE) KOFOID and CAMPBELL.

Text-figure 12.

Text-fig. 10. *Tintinnopsis radix* (IMHOFF) $\times 250$.

Codonella pusilla CLEVE, * 1900; BRANDT, 1907, p. 120.

Codonelllopsis pusilla, Kofoid and CAMPBELL, 1929, p. 87, fig. 146.

Lorica stout fusiform, 2.6 oral diameters in length; collar subcylindrical in the anterior 0.6 of its length and posteriorly conical (55°), then gradually changing into an ovoidal bowl, with about 13 spiral turns extending towards the bowl; bowl ovate, 1.8 of the oral diameter in transdiameter; aboral region an inverted cone of 125° , with a blunt aboral end; wall nearly uniform in thickness throughout the lorica, composed of an alveolar structure and very few agglomerated particles.

Length, $56\ \mu$; oral diameter, $21\ \mu$; transdiameter, $37\ \mu$.

Occurs in July; very rare.

Differs from *Codonelllopsis contracta* Kofoid and CAMPBELL in lack of fenestration of the basal portion of the collar and in the subacute aboral end.

16. *Codonelllopsis contracta* Kofoid and CAMPBELL.


Text-figure 13.

Codonelllopsis contracta Kofoid and CAMPBELL, 1929, p. 78, fig. 147.

Lorica ovate, 2.1 oral diameters in length; collar 0.31–0.35 of the total length in length, usually with 6 spiral turns, concave conical (30° – 40°), provided with a few, transversely elliptical fenestrae in its basal part; bowl globose, 1.7 of the oral diameter in thickness, with primary and secondary structures and few agglomerated particles.

Length, $43(40\text{--}45)\ \mu$; oral diameter, $22(20\text{--}23)\ \mu$; transdiameter, $36(35\text{--}37)\ \mu$.

Occurs in July; rare.



Text-fig. 13. *Codonelllopsis contracta* Kofoid and CAMPBELL. $\times 750$.

Text-fig. 11. *Stenosemella nivalis* (MEUNIER) $\times 750$.

Text-fig. 12. *Codonelllopsis pusilla* (CLEVE) $\times 750$.

Differs from *Codonelllopsis frigida* HADA in having fenestrae and in the smoothly sloping shoulder and from *C. pusilla* (CLEVE) in the presence

of fenestration at the lower part of the collar, in the comparatively small number of spiral turns, and in its hemispherical aboral end.

17. *Codonelllopsis limosa*, n. sp.

Text-figure 14.

Lorica ovoidal, 2.4 oral diameters in length; collar short with a little oral eversion, 0.13 of the total length in length, convex conical (30°), figured with a few spiral turns; bowl ovate, widest near the middle of the lorica, 1.8 oral diameters in transdiameter; aboral region hemispherical; wall agglomerated neatly with small particles.

Length, $82\ \mu$; oral diameter, $33\ \mu$; greatest transdiameter, $63\ \mu$.
Occurs in January; very rare.

Differs from all other allied species of *Codonelllopsis morchella* (CLEVE) in the short collar and in fewer spiral turns.


18. *Codonelllopsis orientalis*, n. sp.

Text-figure 15.

Text-fig. 14. *Codonelllopsis limosa*, n. sp. $\times 350$.



Lorica consisting of a subcylindrical collar and an ovate bowl, 2.5–2.7 oral diameters in length; oral rim flaring; collar 0.30–0.42 of the total



Text-fig. 15. *Codonelllopsis orientalis*, n. sp. $\times 650$.

length in length, constricted slightly in the suboral part and bulging out in the middle, with 8-11 spiral turns increasing in width towards the bowl, few fenestrae appearing in the dilated region; bowl widest a little above the middle, contracting aborally; aboral region broadly convex conical (85° - 105°); aboral end blunt; wall coarsely agglomerated.

Length, $90(84\text{-}92)\mu$; oral diameter, $34(32\text{-}34)\mu$; greatest transdiameter, 50μ .
Occurs in September-January; common.
Differs from *Codonellopsis americana* KOFOID and CAMPBELL, *C. erytreaensis* (BRANDT), and *C. indica* KOFOID and CAMPBELL in the conical aboral region and from *C. morechella* (CLEVE) in an everted oral margin and in the shape of the collar.

Family Cyttarocylidae.
Genus CYTTAROCYLIS Fol, 1881.
19. *Cyttarocylis magna* BRANDT
Text-figure 16.

Cyttarocylis cassis var. *c. magna* BRANDT, 1906, pl. 34, fig. 3, pl. 35, fig. 3; 1907, pp. 189, 196; LACKMANN, 1909, p. 447.
Cyttarocylis magna, KOFOID and CAMPBELL, 1929, p. 114, fig. 222; KOFOID, 1930, fig. 29 (No. 222).

Lorica consisting of a low funnel-shaped collar and a tall inverted conical bowl of angles changing from 15° in the anterior half to 45° in the aboral region, its length 1.8-2.2 oral diameters; oral rim irregularly dentate; collar convex conical (42°), 0.9 oral diameters in transdiameter, 0.06 of the total length in length; aboral horn short, 0.05 of the total length in length, shaped irregularly; wall with a comparatively larger polygonal reticulation.
Length, 300μ ; oral diameter, 150μ .
Occurs in November; rare.

Differs from *Cyttarocylis acutiformis* KOFOID and CAMPBELL in the coarsely reticulated wall and in having the marked aboral horn and from *C. cassis* (HAECKEL) in the tall tapering bowl.

Genus PARAFAVELLA KOFOID and CAMPBELL, 1929.

20. *Parafavella denticulata* (EHRENBURG) KOFOID and CAMPBELL.

Tintinnus denticulatus EHRENBURG, * 1840.
Parafavella denticulata, KOFOID and CAMPBELL, 1929, p. 163, fig. 310; HADA, 1932 b, p. 50, fig. 15.

Length, $225(200\text{-}235)\mu$; oral diameter, $63(62\text{-}64)\mu$.
Occurs in January-March; common.

21. *Parafavella gigantea* (BRANDT) KOFOID and CAMPBELL.
Cyttarocylis gigantea (part) BRANDT, 1896, p. 63, pl. 3, figs. 21, 24.
Parafavella gigantea, KOFOID and CAMPBELL, 1929, p. 165, fig. 311; HADA, 1932 b, p. 51, fig. 16.


Length, $380(337\text{-}486)\mu$; oral diameter, $65(63\text{-}69)\mu$.
Occurs in January-May; common.

22. *Parafavella pacifica* HADA.
Parafavella pacifica HADA, 1932 b, p. 49, fig. 13.

Length, $135(120\text{-}154)\mu$; oral diameter, $47(45\text{-}52)\mu$.
Occurs in June-September; common.

23. *Parafavella faceta*, n. sp.
Text-figure 17

Lorica tall bell-shaped, 2.0-3.1 oral diameters in length; oral rim denticulate with about 48 triangular teeth; bowl dilated slightly at the



Text-fig. 16. *Cyttarocylis magna* BRANDT $\times 200$.

Text-fig. 17. *Parafavella faceta*, n. sp. $\times 400$.

Text-fig. 18. *Parafavella venusta* (JÖRGENSEN) $\times 250$.

Y. HADA

Family *Ptychocylidae*.Genus *PTYCHOCYLIS* BRANDT, 1896.26. *Ptychocylis obtusa* BRANDT.

Posterior 0.2 of the total length, its widest transdiameter 1.07 oral diameters, below the suboral expansion contracting gradually, changing from 10° – 15° at the middle to 70° – 90° in the aboral region; aboral horn stout, shortened, conical (20° – 32°), 0.07–0.09 of the total length in length, tip usually acute. Length, $147(123$ – $171)\mu$; oral diameter, $56(51$ – $61)\mu$; greatest transdiameter, $64(54$ – $65)\mu$.

Occurs in January; common.

Differs from *Parafavella obtusangula* (OSTENFELD) and *P. parvudentata* (BRANDT) in more abrupt contraction of the aboral region and in having many more distinct aboral horn and from *P. pacifica* HADA in having many teeth of the oral rim and in the slowly contracting aboral part.

24. *Parafavella ventricosa* (JÖRGENSEN) KOFOID and CAMPBELL. Text-figure 18.

Text-figure 18.

Cyathocylis denticulata var. β *cylindrica* forma *ventricosa* JÖRGENSEN, 1899, p. 34, pl. 3, fig. 30.

Parafavella ventricosa, KOFOID and CAMPBELL, 1929, p. 171, fig. 314; KOFOID, 1930, fig. 30 (No. 314).

Lorica finger-shaped, 5 oral diameters in length; oral margin denticulated with comparatively fewer teeth; bowl subcylindrical, with a slight suboral bulge, enlarged gradually towards the aboral end and widest at the posterior aboral region an inverted convex cone of 78° ; aboral horn 0.1 of the total length in length, conical (8°), tip more or less acute.

Length, 316μ ; oral diameter, 64μ ; greatest transdiameter, 80μ .

Occurs in August; very rare.

Differs from *Parafavella gigantea* (BRANDT) in the bulbous aboral region, in the shorter horn, and in fewer teeth on the oral margin.

25. *Parafavella subrotunda* (JÖRGENSEN) KOFOID and CAMPBELL.

Text-figure 19.
Genus *ACANTHOSTOMELLA* JÖRGENSEN, 1927.

Cyathocylis denticulata var. γ *subrotunda* JÖRGENSEN, 1899, p. 34, pl. 2, figs. 20, 21. *Porefavella subrotunda*, KOFOID and CAMPBELL, 1929, p. 170, fig. 316; HADA, 1932 b, p. 54, fig. 1.

Length, $220(211$ – $228)\mu$; oral diameter, $62(58$ – $63)\mu$. Occurs in February and March; rare.

Ptychocylis drygalskii BRANDT, 1896, p. 59, pl. 3, fig. 14; KOFOID and CAMPBELL, 1929, p. 188, fig. 350.
Ptychocylis urmala var. *digitalis* JÖRGENSEN, 1901, p. 17, pl. 2, figs. 29, 30.
Ptychocylis urmala var. *digitalis* forma *subintegerrima* JÖRGENSEN, 1901, p. 26, pl. 3, fig. 31.

Ptychocylis obtusa var. *drygalskii* (part), BRANDT, 1906, pl. 55, figs. 1–3, pl. 56, figs. 3, 3 a, pl. 57, fig. 10; 1907, p. 312.

Ptychocylis ventricosa MEUNIER, 1910, p. 127, pl. 10, fig. 3.
Lorica wide goblet-shaped, 1.4 oral diameters in length; oral rim regularly denticulated; bowl generally a low, inverted, convex cone with two distinct expansions: each respectively 1.15 and 1.20 oral diameters in transdiameter; aboral region concave conical 90° ; aboral end broadly rounded, more or less thicker than the other parts, with a rugose surface.

Length, 77μ ; oral diameter, 55μ .
Occurs in July; very rare.
Differs from *Ptychocylis obtusa* BRANDT in stouter proportions and in the shape of the aboral cone.

27. *Ptychocylis drygalskii* BRANDT.
Text-figure 19.

Text-figure 19.

Amphorella norvegica DADAY, 1887, p. 543.
Acanthostomella norvegica, KOFOID and CAMPBELL, 1929, p. 193, fig. 363; HADA, 1932 b, p. 56, fig. 22.

Length, 43 – 47μ ; oral diameter, 29 – 30μ ; greatest transdiameter, 31 – 32μ .
Occurs in February and March; rare.

Family Xystonellidae.

Genus PARUNDELLA JÖRGENSEN, 1924.

29. *Parundella pellucida* (JÖRGENSEN) KOFOID and CAMPBELL.

Text-figure 21.

Undella pellucida (part) JÖRGENSEN, 1899, p. 41, pl. 1, fig. 7.


Parundella pellucida, KOFOID and CAMPBELL, 1929, p. 233, fig. 438.

Lorica elongate chalkite-shaped, 4 oral diameters in length; oral rim entire; bowl subcylindrical, slightly contracting in the anterior 0.25 of the oral length; aboral region tapering (45°) to a caudal spine; lance conical (18°), 0.25 of the total length in length, single-lamellate in the posterior 0.42 of its length, with three somewhat spiral costae on the upper thick-walled part, tip pointed.

Length, $102\ \mu$; oral diameter, $27\ \mu$.

Occurs in November; very rare.

Differs from *Parundella caudata* (OSTENFELD) in the absence of predominant fins at the junction of the aboral region and the lance.



Text-fig. 19 *Psychoclysis drygalskii* BRAUDER $\times 600$.

Text-fig. 20. *Acanthostomella norvegica* (DADAY) $\times 1000$.

Text-fig. 21. *Parundella pellucida* (JÖRGENSEN) $\times 600$.

Family Undellidae.


Genus PROPLECTELLA KOFOID and CAMPBELL, 1929.

30. *Proplectella expolita*, n. sp.

Text-figure 22.

Lorica stout flask-shaped, with a well-developed outer collar, 2.00–2.58

oral diameters in length; suboral region subcylindrical, thinning outwardly to a sharp oral rim, its length 0.30–0.33 of the total length; bowl generally subspherical, sometimes broadly ovate, 1.5–2.0 oral diameters



Text-fig. 22. *Proplectella expolita*, n. sp. $\times 750$.

in transdiameter; aboral end usually hemispherical or widely rounded, occasionally showing weakly a trace of bluntly pointing; wall thickened at the basal part of the collar, thinning gradually towards the aboral end, about 0.15 of the oral diameter in thickness in the thickest wall of the collar.

Length, $66(56\text{--}76)\ \mu$; oral diameter, $27(24\text{--}30)\ \mu$; transdiameter of the bowl, $49(44\text{--}52)\ \mu$.

Occurs in June–August; common.

Differs from all other species of *Proplectella* in having the conspicuous collar and from *Undella californiensis* KOFOID and CAMPBELL in the structure of the wall which is uniform in thickness in that species, but in this species is thickest at the lower part of the collar and thinnest at the aboral end.

Family Tintinnidae.

Genus AMPHORELLA DADAY, 1887.

31. *Amphorella brandti* JÖRGENSEN.

Text-figure 23.

Tintinnus amphora, BRAUDER, 1906, pl. 69, fig. 6; 1907 (part), p. 433.
Amphorella quadrilineata var. *brandti* JÖRGENSEN, *1924.

Amphorella brandti, KOFOID and CAMPBELL, 1929, p. 309, fig. 588.

Lorica 2.2–2.6 oral diameters in length, consisting of a collar of the circular cross-section and a triangular bowl, narrowest at the base of the

33. *Tintinnus tenuis* KOFOID and CAMPBELL.

Text-figure 25.

Tintinnus lasus-undae (part) ZACHARIAS, 1906, p. 518, fig. 6; BRANDT, 1907, p. 420.

Tintinnus lasus-undae var. c (part), BRANDT, 1906, pl. 65, fig. 10; 1907, p. 422.

Tintinnus tenuis KOFOID and CAMPBELL, 1929, p. 339, fig. 655.

Lorica a long, inverted, truncated cone of 2° - 3° , with very slight expansion in the middle part, its length 4.1-5.9 oral diameters; both end entire, oral one flaring with a brim, the other without a flare, 0.61-0.71 of the oral diameter in aboral diameter. Length, 261(236-312) μ ; oral diameter, 51(49-56) μ .

Occurs in March-October; common.

Differs from *Amphorella quadrilineata* (CLAPARÈDE and LACHMANN) in having three instead of four fins.

Genus TINTINNUS SCHRANK, 1803.

32. *Tintinnus exigua*, n. sp.



Text-figure 24.

Lorica an elongate truncated cone of 3° - 5° , 3.5-4.2 oral diameters in length; oral end abruptly flaring to form a marked brim; sides nearly straight; aboral end 0.64-0.75 of the oral diameter in aboral diameter, without a brim and a flare.

Length, 148(130-160) μ ; oral diameter, 38(36-40) μ .

Occurs in May-October; common.

Differs from *Tintinnus tenuis* KOFOID and CAMPBELL in being smaller and more widely conical.



Genus SALPINGELLA JÖRGENSEN, 1921.

34. *Salpingella attenuata* JÖRGENSEN.

Text-figure 26.

Tintinnus acuminatus, ENTZ, Sr., 1885, p. 201, pl. 14, fig. 13.

Tintinnus acuminatus var. c *gluckentoweni* (part) BRANDT, 1906, pl. 68, fig. 2-4; 1907, p. 330.

Salpingella acuminata subsp. *gluckentoweni* var. *attenuata* JÖRGENSEN, 1921.

Salpingella attenuata, KOFOID and CAMPBELL, 1929, p. 351, fig. 687.

Lorica much elongated, slender, 9 oral diameters in length; consisting of a funnel-shaped collar and a long tubular bowl; collar a wide inverted concave cone of 60° ; bowl cylindrical in the posterior half, 0.5 of the oral diameter in length, sharply conical (5°) in the posterior half, distally convex conical (30°); aboral end open, truncate; aboral region with 6 somewhat dextro-tropic fins on the posterior 0.29 of the total length.

Length, 288 μ ; oral diameter, 32 μ ; trans-diameter of the bowl, 16 μ .

Occurs in November; very rare.

Differs from *Salpingella gracilis* KOFOID and CAMPBELL in size and in having fewer fins and

Text-fig. 23. *Amphorella brandti* JÖRGENSEN $\times 500$.

Text-fig. 24. *Tintinnus exigua*, n. sp. $\times 400$.

Text-fig. 25. *Tintinnus tenuis* KOFOID and CAMPBELL $\times 250$.

Text-fig. 26. *Salpingella attenuata* JÖRGENSEN $\times 300$.

TINTINNOINEA OF MUTSU BAY

from *S. ricta* KOFOID and CAMPBELL in more slender proportions and in lack of the surface rugose.

BIBLIOGRAPHY

- ECKARDT, K., 1896. Zoologische Ergebnisse der von der Gesellschaft für Erdkunde zu Berlin unter Leitung Dr. von DRAGALSKI's ausgesandten Grönlandexpedition nach Dr. VANHOFFEN's Sammlungen bearbeitet. IV. Die Tintinen.—Bibliotheca Zoologica, Bd. 8, pp. 45-72, pl. 3.
- 1906. Die Tintinnoden der Plankton-Expedition. Tafelklärungen nebst kurzer Diagnose der neuen Arten.—Ergebn. Plankton-Exped. Humboldt-Stift., Bd. 3, L. a., 33 pp., 70 pls.
- 1907. Die Tintinnoden der Plankton-Expedition. Systematischer Theil.—Ergebn. Plankton-Exped. Humboldt-Stift., Bd. 3, L. a., 488 pp.
- CAMPBELL, A. S., 1926. The Cytology of *Tintinnopsis mucula* (FOL) LAACKMANN with an Account of its Neuromotor Apparatus, Division, and a new Intracellular Parasite. Univ. California Publ. Zool., Vol. 29, pp. 179-236, pls. 12-15, 7 text-figs.
- CLEVE, P. T., 1899. Plankton collected by the Swedish Expedition to Spitzbergen in 1898.—Kgl. Svenska Vet.-Akad Handl., Bd. 32, No. 3, 51 pp., 4 pls.
- 1900 a. The Plankton of the North Sea, the English Channel, and the Skagerak in 1898.—Kgl. Svenska Vet.-Akad Handl., Bd. 32, No. 8, 53 pp., 11 text-figs.
- 1900 b. The Plankton of the North Sea, the English Channel and the Skagerak in 1899.—Kgl. Svenska Vet.-Akad Handl., Bd. 34, No. 2, 77 pp.
- 1902. The Plankton of the North Sea and the Skagerak in 1900.—Kgl. Svenska Vet.-Akad. Handl., Bd. 35, No. 7, 49 pp., 1 text-fig.
- 1903. Plankton-researches in 1901 and 1902.—Kgl. Svenska Vet.-Akad. Handl., Bd. 36, No. 8, 53 pp., 2 text-figs.
- DABAY, E. v., 1886. Ein kleiner Beitrag zur Kenntnis der Infusorien-Fauna des Golfs von Neapel.—Mitt. Zool. Sta. Neapel, Bd. 6, pp. 481-498, pl. 25.
- 1887. Monographie der Familie der Tintinnoden.—Mitt. Zool. Sta. Neapel, Bd. 7, pp. 473-591, pls. 18-21.
- ENTZ, G. Sr., 1884. Über Infusorien des Golfs von Neapel.—Mitt. Zool. Sta. Neapel, Bd. 5, pp. 289-444, pls. 20-25.
- 1885. Zur näheren Kenntnis der Tintinnoden.—Mitt. Zool. Sta. Neapel, Bd. 6, pp. 185-216, pls. 13, 14.
- ENTZ, G. Jr., 1909. Studien über Organisation und Biologie der Tintinniden.—Arch. Prot., Bd. 15, pp. 93-226, pls. 8-21, 2 text-figs.
- FOL, H., 1881. Contribution to the Knowledge of the Family Tintinnidae.—Ann. Mag. Nat. Hist., Ser. 5, Vol. 7, pp. 237-250, pl. 17, figs. 1-6.
- HADA, Y., 1932 a. Descriptions of two new neritic Tintinnoinea, *Tintinnopsis japonica* and *Tps. kofoidi* with a brief Note on a Unicellular Organism parasitic on the latter.—Proc. Imp. Acad., Vol. 8, No. 5, pp. 209-212, 3 text-figs.
- 1932 b. The Tintinnoinea from the Sea of Okhotsk and its Neighborhood.—Journ. Fac. Sci. Hokkaido Imp. Univ., Ser. 4, Vol. 2, No. 1, pp. 37-59, 23 text-figs.
- IMHOF, O. E., 1886. Über mikroskopische pelagische Thiere aus den Lagunen von Venedig.—Zool. Anz., Bd. 9, pp. 101-104.
- JÖRGENSEN, E., 1889. Über die Tintinnoidea der norwegischen Westküste.—Berger's Mus.
- 1900. Protozoen und Protozoen im Plankton aus der norwegischen Westküste.—Bergens Mus. Arbog, 1899, No. 6, 112+LXXXIII pp., 5 pls.
- 1901. Protistenplankton aus dem Nordmeere in den Jahren 1897-1900.—Bergens Mus. Arbog, 1900, No. 6, 37 pp., 3 pls.
- 1927. "Ciliata: Tintinnidae".—Die Tierwelt der Nord- und Ostsee, Lief. VIII, Teil II. C, 26 pp., 33 text-figs.
- KENT, W. S., 1881-1882. A Manual of the Infusoria (Tintinnoinea, 1882). 3 vols., X-913 pp., 51 pls.
- KOFOID, C. A., 1905. Some New Tintinnidae from the Plankton of the San Diego Region.—Univ. California Publ. Zool., Vol. 1, pp. 287-306, pls. 26-28.
- 1930. Factors in the Evolution of the Pelagic Ciliata, the Tintinnoinea.—Cont. Mar. Biol. Stanford Univ., 39 pp., 1 pl., 31 text-figs.
- KOFOID, C. A. and A. S. CAMPBELL, 1929. A Conspectus of the Marine and Fresh-water Ciliata belonging to the Suborder Tintinnoinea, with Descriptions of New Species principally from the Agassiz Expedition to the Eastern Tropical Pacific 1904-1905.—Univ. California Publ. Zool., Vol. 34, 403 pp., 697 text-figs.
- LAACKMANN, H., 1906. Ungeschlechtliche und geschlechtliche Fortpflanzung der Tintinen.—Wiss. Meeresunters., N. F., Bd. 10, pp. 15-38, pls. 1-3.
- 1913. Adriatische Tintinnoidea.—Sitzungsber. Kais. Akad. Wiss. Wien. Mathe-nat. Klasse, Bd. 122, pp. 1-45, pls. 1-6, 2 text-figs.
- MERESCHKOWSKY, C., 1879. Studien über Prozoen des nördlichen Russland.—Arch. Mikr. Anat., Bd. 16, pp. 153-248, pls. 10, 11.
- MERKELE, H., 1909. Untersuchungen an Tintinnoidea der Ost- und Nordsee.—Wiss. Meeresunters., Bd. 11, pp. 139-186, pls. 2, 3, 3 text-figs.
- MEUNIER, A., 1910. Micoplankton des Mers de Barent et de Kara.—Duc D'ORLEANS, Comm. pagne Arctique de 1907, XVIII+355 pp., 37 pls.
- OKAMURA, K., 1907. An annotated List of Plankton Microorganisms of the Japanese Coast.—Annal. Zool. Japoneses, 1906-1908, Vol. 6, pp. 125-151, pls. 3-6.
- ZACHARIAS, O., 1906. Über Periodicität, Variation und Verbreitung verschiedener Planktonwesen in südlichen Meeren.—Arch. Hydrobiol. Plankton., Bd. 1, pp. 428-575, pl. 1, 23 text-figs.